

Psoas Versus Femoral Blocks

A Registry Analysis of Risks and Benefits

Hagen Bomberg, MD,* Andrea Huth, MS,* Stefan Wagenpfeil, PhD,† Paul Kessler, MD,‡ Hinnerk Wulf, MD,§ Thomas Standl, MD,|| André Gottschalk, MD,** Jens Döffert, MD,†† Werner Hering, MD,‡‡ Jürgen Birnbaum, MD,§§ Claudia Spies, MD,§§ Bernd Kutter, MD,|||| Jörg Winkelmann, MD,||||| Gerald Burgard, MD,*** Oliver Vicent, MD,††† Thea Koch, MD,††† Daniel I. Sessler, MD,‡‡‡ Thomas Volk, MD,* and Alexander Raddatz, MD*

Background and Objectives: Psoas blocks are an alternative to femoral nerve blocks and have the potential advantage of blocking the entire lumbar plexus. However, the psoas muscle is located deeply, making psoas blocks more difficult than femoral blocks. In contrast, while femoral blocks are generally easy to perform, the inguinal region is prone to infection. We thus tested the hypothesis that psoas blocks are associated with more insertion-related complications than femoral blocks but have fewer catheter-related infections.

Methods: We extracted 22,434 surgical cases from the German Network for Regional Anesthesia registry (2007–2014) and grouped cases as psoas ($n = 7593$) and femoral ($n = 14,841$) blocks. Insertion-related complications (including single-shot blocks and catheter) and infectious complications (including only catheter) in each group were compared with χ^2 tests. The groups were compared with multivariable logistic models, adjusted for potential confounding factors.

Results: After adjustment for potential confounding factors, psoas blocks were associated with more complications than femoral blocks including vascular puncture 6.3% versus 1.1%, with an adjusted odds ratio (aOR) of 3.6 (95% confidence interval [CI], 2.9–4.6; $P < 0.001$), and multiple skin punctures 12.6% versus 7.7%, with an aOR of 2.6 (95% CI, 2.1–3.3; $P < 0.001$). Psoas blocks were also associated with fewer catheter-related infections: 0.3% versus 0.9% (aOR of 0.4; 95% CI, 0.2–0.8; $P = 0.016$), and with improved patient satisfaction (mean \pm SD 0- to 10-point scale score,

9.6 ± 1.2 vs 8.4 ± 2.9 ; $P < 0.001$). Results from a propensity-matched sensitivity analysis were similar.

Conclusions: Psoas blocks are associated with more insertion-related complications but fewer infectious complications.

Clinical Trial Registration: ID NCT02846610.

(*Reg Anesth Pain Med* 2017;42: 719–724)

Various peripheral nerve blocks can be used for major hip and knee surgery. In 1976, Chayen et al described the psoas block as a posterior approach to the lumbar plexus.¹ Since then, several works refining this regional anesthetic technique have been published.²

Severe and even life-threatening hemorrhagic complications have been described after psoas blocks with or without catheter insertion.^{2–4}

Consequently, the American Society of Regional Anesthesia and Pain Medicine recommends applying the same guidelines for this “deep” peripheral nerve block as for neuraxial blocks in patients given anticoagulant or antiplatelet therapy.⁵ Local anesthesia systemic toxicity after unintended intravascular injection has also been described.⁶ Other serious adverse effects include cardiac arrest, respiratory failure, and seizures related to cephalad diffusion of the local anesthetic in the epidural or intrathecal spaces.^{7,8}

Femoral nerve blocks are an alternative to psoas blocks and are considered among the easiest-to-perform peripheral blocks. Although systemic local anesthesia toxicity including cardiac arrest can occur after any block, the incidence is believed to be lower after femoral than psoas blocks.^{8–10} However, femoral vein catheter placement has been shown to be associated with a higher rate of bloodstream infections compared with upper-body central venous vascular catheter placement.¹¹

Available evidence thus suggests that insertion-related complications are more likely with psoas blocks, but that catheter-related infections may be less common. We therefore compared the incidences of complications during insertion and catheter-related infections associated with psoas and femoral blocks in a large registry. Specifically, we evaluated the hypotheses that insertion-related complications for single-injection blocks and catheter are more common with psoas than femoral blocks and that catheter-related infections are less common.

METHODS

This trial was registered under Ha50/11 and approved by the ethics committee of the Saarland Medical Chamber, Saarbrücken, Germany (Chairperson San.-Rat Prof Dr Hermann Schieffer) on March 22, 2011. According to the regulatory proof of protection of data privacy (Saarland commissioner, 12-MAR-2014), written consent was waived as the data were anonymous. This trial was also registered at clinicaltrials.gov on May 28, 2016 (identifier NCT02846610).

From the *Department of Anesthesiology, Intensive Care Medicine and Pain Medicine, and †Institute for Medical Biometry, Epidemiology and Medical Informatics, Saarland University, University Medical Center, Homburg/Saar; ‡Department of Anesthesiology, Intensive Care and Pain Medicine, Orthopedic University, Hospital, Frankfurt; §Department of Anesthesiology and Intensive Care Therapy, Philipps University Marburg, Marburg; ||Department of Anesthesia, Intensive and Palliative Care Medicine, Solingen Academic Hospital of the University of Cologne, Solingen; **Department of Anesthesiology, Intensive Care and Pain Medicine, Friederikenstift Hospital, Hannover; ††Department of Anesthesiology and Intensive Care Medicine, Hospital Calw-Nagold; ‡‡Department of Anesthesiology, St Marien-Hospital, Siegen; §§Department of Anesthesiology and Operative Intensive Care Medicine, Charité Campus Virchow Klinikum and Campus Mitte, Charité University Medicine, Berlin; ||||Department of Anesthesiology, Intensive Care and Pain Therapy, University and Rehabilitation Clinics, Ulm; ***Department of Anesthesiology, Intensive Care and Pain Therapy, HELIOS Hospital Erfurt, Erfurt; †††Department of Anesthesiology, Intensive Care and Pain Medicine, University Hospital, Dresden, Germany; and ‡‡‡Department of Outcomes Research, Anesthesiology Institute, Cleveland Clinic, Cleveland, OH.

Accepted for publication March 22, 2017.

Address correspondence to: Hagen Bomberg, MD, Department of Anesthesiology, Intensive Care Medicine and Pain Medicine, Saarland University Medical Center, Kirrbergerstrasse 1, 66421 Homburg/Saar, Germany (e-mail: Hagen.Bomberg@uniklinikum-saarland.de).

This analysis was supported by institutional funds. The German Network for Regional Anesthesia was supported by the German Society of Anesthesiology and Intensive Care Medicine, the Association of German Anesthesiologists, and the German Research Foundation.

The authors declare no conflict of interest.

Copyright © 2017 by American Society of Regional Anesthesia and Pain Medicine

ISSN: 1098-7339

DOI: 10.1097/AAP.0000000000000643

The German Network for Regional Anesthesia, which was established in 2007 by the German Society for Anesthesiology and Intensive Care Medicine and the Professional Association of German Anesthesiologists, set up a database that collects preoperative, intraoperative, and postoperative data from treating physicians or pain nurses at 25 German centers using a standard form (Fig. 1).¹² Data were collected concurrently with patient care and included detailed information about the medical conditions of patients having regional anesthesia along with details of the procedure and postoperative course.

The registry investigated 126,236 cases in the period extending from September 2007 to May 2015. According to specific rules to delete erroneously entered data and to delete cases with missing information, data integrity was evaluated, and the relation between age, height, weight, and sex were verified. The body mass index (BMI) was defined from 13.2 to 70 kg/m². Anticoagulant therapy comprised warfarin and platelet inhibition, as well as high- and low-molecular-weight heparin.

All participating centers administered routine single-dose antibiotic prophylaxis based on the surgical procedure and recommended by the guidelines of the German working group of Hygiene in Hospital and Practice.^{13,14} They complied with the German guidelines to prevent catheter-related infection and block-related hemorrhage in patients receiving antithrombotic or thrombolytic therapy or having coagulation disorders.^{15,16}

Case Selection

We included patients who received femoral or psoas blocks with or without concomitant sciatic nerve block. Exclusion criteria were catheter use exceeding 14 days, patients with missing information about age and sex, patients with diabetes, and patients with implausible data. Implausible data included a diagnosis of diabetes and an American Society of Anesthesiologists (ASA) physical status score of 1 or the date of block insertion preceding the date of birth.

Block-related complications and patient satisfaction were prospectively defined and included the following:

- 1. vascular puncture: aspiration of blood by the puncture needle or catheter;
- 2. multiple skin punctures, defined by more than 1 skin puncture;
- 3. paresthesia during needle or catheter insertion: an unexpected painful, unpleasant, or electric sensation within the area innervated by the nerve(s) affected by the regional anesthesia;
- 4. accidental dural puncture: unintentional needle insertion through the dura mater;
- 5. systemic local anesthetic toxicity: symptoms of intoxication (neurologic and/or cardiovascular) after the injection of the local anesthetic;

- 6. premature termination (of the block procedure): procedure terminated because of unfavorable anatomic conditions, insufficient compliance of the patient, or vasovagal symptoms during the procedure; and
- 7. primary failure: the block did not provide intraoperative analgesia.
- 8. Infections at the catheter insertion site were categorized by severity: (a) mild infections were defined by at least 2 of 3 infection signs (redness, swelling, or local pain); (b) moderate infections were defined as mild plus by at least 1 of the following findings: increased C-reactive protein, leukocytosis, fever, or pus at the punctured site; and (c) severe infections were defined by the need for a surgical incision or revision.
- 9. Patients and provider satisfaction were measured with verbal numeric rating scales ranging from 0 (completely dissatisfied) to 10 (completely satisfied) at the last encounter.

Data Analysis

Population characteristics are reported as standardized differences (psoas – femoral). Dichotomous outcomes in patients with psoas and femoral block were compared with χ^2 tests. Continuous variables were compared with Student *t* tests (respectively, Welch *t* tests in case of unequal variances).

Logistic regression analysis was used to calculate univariate and adjusted odds ratios (aORs) with 95% confidence intervals (CIs). We adjusted for factors known to influence block-related complications including ASA physical status, diabetes, anticoagulant therapy, increased BMI, and type of block.^{9,10,17,18} Additional relevant confounders were identified for each block-related complication (vascular puncture, multiple skin punctures, infection for any grade) and for satisfaction of patients and physician. Finally, we included year of surgery and hospital. Collinearity was tested by Pearson or Spearman correlation coefficients. Variables with a positive or negative correlation greater than +0.3 and less than –0.3 were assessed for interactions. Goodness of fit was assessed by Hosmer-Lemeshow tests.

A propensity score was developed for each patient based on all potential confounders and implemented as a sensitivity analysis. The patient with the closest propensity score who had a femoral block was matched to each patient with a psoas block. The maximum difference of propensity score for a matching was less than 0.05. Matched patients with psoas and femoral block were compared using McNemar test, Student *t* tests for 2 dependent samples, or conditional logistic regression. Covariable balance for propensity score matching was assessed with standardized differences. Imbalance was defined as a standardized difference of greater than 0.1 in absolute value; any imbalanced covariables

Preoperative data	Block placement and intraoperative data	Postoperative data and ward round
Male	Catheter / Single shot	Infection
Age [yrs.]	Site	Length of catheter use
Body mass index [kg/m ²]	Antibiotic prophylaxis	Patient satisfaction with RA
ASA physical status	Nerve stimulation	Physician satisfaction with RA
Diabetes mellitus	Ultrasound	
Anticoagulant therapy	Vascular puncture	
Department of surgery	Multiple skin puncture	
	Paresthesia	
	Accidental dural puncture	
	Local anesthetic systemic toxicity	
	Premature termination anatomy	
	Premature termination compliance	
	Primary failure	

FIGURE 1. Extract of the information completed after block placement and ward rounds. RA indicates regional anesthesia.

after the propensity score matching were adjusted in a further sensitivity analysis.

We used Python Essentials, an extension of SPSS, for propensity score matching. All data analyses were performed using IBM SPSS Statistics for Windows (version 22; IBM Corp, Armonk, New York). Continuous variables are expressed as means \pm SDs. Categorical variables are presented as numbers of patients with percentages in parentheses. Statistical significance was accepted at a global 2-sided significance level of 0.05. Because of multiple testing, we considered Bonferroni correction to arrive at a local significance level of 0.05/9 (primary outcomes) = 0.006.

RESULTS

The final study population consisted of 22,434 cases: all 7593 qualifying psoas blocks and 14,841 femoral blocks (Fig. 2).

Patients with psoas blocks were more often female, were older, had a higher (ASA) physical status score, used more anticoagulant therapy, and were more often given antibiotic prophylaxis. A greater fraction of psoas block patients had catheters inserted, and psoas blocks were more often guided by nerve stimulation (Table 1).

Patients who had psoas blocks had more vascular punctures, more often needed multiple punctures, were more likely to experience systemic toxicity of local anesthetics, and more often had premature termination of the insertion attempt. In contrast, there were fewer cases of paresthesia, primary failures, and block-related infections in patients with psoas block. Catheter duration was longer, and the satisfaction of patients and providers was higher in patients with psoas block compared with femoral block (Table 2). No cardiac arrests were reported consequent to systemic local anesthetic toxicity. No deaths were attributed to either psoas or femoral blocks.

After multivariable analysis, adjusted for potential confounding factors, psoas blocks were associated with increased risk of vascular punctures and multiple skin punctures compared with femoral

blocks. In contrast, psoas blocks were associated with fewer primary failures, fewer block-related infections, and more satisfied patients and providers both before and after adjustment for confounders (Fig. 3). Goodness of fit for each adjusted model was assessed by Hosmer-Lemeshow tests and did not differ significantly.

Propensity matching successfully paired 5928 patients with psoas block (78% of 7593 patients) and with 5928 control subjects (40% of 14,841 patients). As seen in Table 1, matched patients with psoas and femoral blocks were much better balanced on covariables. However, imbalances remained for BMI (standardized difference [STD], -0.16), antibiotic prophylaxis (STD, 0.14), type of block (STD, 0.19), use of nerve stimulator (STD, 0.13), use of ultrasound (STD, -1.20), year of surgery (STD, -0.15), and hospital (STD, 0.50). To be conservative, we included each unbalanced factor in our multivariable model. In addition, satisfaction of patients and providers was also adjusted for vascular punctures, multiple punctures, primary failure, block-related infection, and duration of catheter use. The results were thus similar to those obtained from our unmatched multivariable analysis (Fig. 3).

DISCUSSION

Although major block-related complications such as systemic local anesthetic toxicity and infection are rare, they are potentially serious. Few, if any, previous studies had sufficient capacity to detect clinically important differences between psoas and femoral blocks or to accurately characterize the risks of either. Our multicenter registry analysis of 22,434 patients appears to be the world's largest systematic collection of psoas and femoral blocks. We were thus able to detect complications, even rare ones.

Psoas blocks were associated with more vascular punctures and more skin punctures than femoral blocks. But psoas blocks were also strongly associated with fewer block-related infections and greater patient satisfaction. Results were similar in our propensity-matched




FIGURE 2. Case selection.

TABLE 1. Population Characteristics

	Before Matching			After Matching		
	Psoas Block	Femoral Block	STD*	Psoas Block	Femoral Block	STD*
	(n = 7593)	(n = 14,841)		(n = 5928)	(n = 5928)	
Male	3245 (42.7)	6619 (44.6)	−0.04	2590 (43.7)	2566 (43.3)	0.01
Age, y	62 ± 19	56 ± 20	0.28	60 ± 20	60 ± 18	0.00
Body mass index, kg/m ²	28.5 ± 5.9	28.2 ± 6.1	0.05	28.2 ± 5.9	29.1 ± 6.0	−0.16
ASA physical status score ≥2	6288 (83)	9595 (65)	0.42	4765 (80)	4879 (82)	−0.05
Diabetes mellitus	1018 (13.4)	1854 (12.5)	0.03	767 (12.9)	762 (12.9)	0.00
Anticoagulant therapy	2890 (38.1)	3967 (26.7)	0.25	2329 (39.3)	2304 (38.9)	0.01
Antibiotic prophylaxis	4814 (63.4)	7419 (50.0)	0.27	4115 (69.4)	3711 (62.6)	0.14
Catheter	6697 (88.2)	11,645 (78.5)	0.26	5106 (86.1)	4672 (78.8)	0.19
Nerve stimulation	6904 (90.9)	8012 (54.0)	0.91	5413 (91.3)	5168 (87.2)	0.13
Ultrasound	369 (4.9)	7578 (51.1)	−1.20	200 (3.4)	2879 (48.6)	−1.20
Sciatic nerve	4011 (52.8)	7596 (51.2)	0.03	2896 (48.9)	2861 (48.3)	0.01
Traumatology and orthopedics	6907 (91.0)	12,329 (83.1)	0.24	5837 (98.5)	5854 (98.8)	−0.02

Continuous variables are expressed as means ± SDs. Categorical variables are presented as numbers of patients with percentages in parentheses.

Other surgery includes general surgery, gynecology, internal medicine, pediatric surgery, urology, and vascular surgery.

Selected variables for pairwise matching were sex, age, BMI, ASA physical status score, diabetes, anticoagulant therapy, antibiotic prophylaxis, catheter, single-shot blocks, nerve stimulation, department of surgery, sciatic nerve catheter (as second catheter), year of surgery, and hospital center.

*The difference (psoas minus femoral) in means or proportions divided by the pooled SD.

TABLE 2. Block-Related Complications, Catheter Duration, and Satisfaction

	Before Matching			After Matching		
	Psoas Block	Femoral Block	P	Psoas Block	Femoral Block	P
	(n = 7593)	(n = 14,841)		(n = 5928)	(n = 5928)	
Vascular puncture	475 (6.3)	170 (1.1)	<0.001	400 (6.7)	107 (1.8)	<0.001
Multiple skin puncture	955 (12.6)	1148 (7.7)	<0.001	759 (12.8)	502 (8.5)	<0.001
Paresthesias during block insertion	31 (0.4)	122 (0.8)	0.001	29 (0.5)	41 (0.7)	0.32
Accidental dural puncture	1 (0.0)	0 (0.0)	0.16	0 (0.0)	0 (0.0)	—
Local anesthetic systemic toxicity	5 (0.1)	0 (0.0)	0.002	4 (0.1)	0 (0.0)	0.045
Premature termination (anatomy)	11 (0.2)	4 (0.0)	0.001	11 (0.2)	1 (0.0)	0.004
Premature termination (compliance)	2 (0.0)	0 (0.0)	0.048	1 (0.0)	0 (0.0)	0.317
Primary failure	57 (0.8)	337 (2.3)	<0.001	16 (0.3)	211 (3.6)	<0.001
Infection						
Any grade	24 (0.3)	134 (0.9)	<0.001	16 (0.3)	42 (0.7)	<0.001
Mild	21 (0.3)	115 (0.8)	<0.001	13 (0.2)	33 (0.6)	<0.001
Moderate	3 (0.0)	16 (0.1)	0.063	3 (0.1)	8 (0.1)	0.071
Severe	0 (0.0)	3 (0.0)	0.19	0 (0.0)	0 (0.0)	—
Missing information	1143 (15.1)	3521 (23.7)	—	1025 (17.3)	1819 (30.7)	—
Prolonged catheter use (4–14 d)	1585 (20.9)	2317 (15.6)	<0.001	1188 (20.0)	642 (10.8)	<0.001
Missing information	1494 (19.7)	4541 (30.6)	—	1327 (22.4)	2314 (39.0)	—
Patient satisfaction (0–10)	9.6 ± 1.2	8.4 ± 2.9	<0.001	9.7 ± 1.1	9.1 ± 1.9	<0.001
Missing information	3202 (42.2)	7609 (51.3)	—	2213 (37.3)	2326 (39.2)	—
Physician satisfaction (0–10)	9.0 ± 2.5	7.7 ± 3.5	<0.001	9.5 ± 1.4	8.8 ± 2.1	<0.001
Missing information	6350 (83.6)	9210 (62.1)	—	4840 (81.7)	3372 (56.9)	—

Continuous variables are expressed as means ± SDs. Categorical variables are presented as numbers of patients with percentages in parentheses. Bold data are significant.

Mild infections were defined by at least 2 out of 3 infection signs (redness, swelling, or local pain). Moderate infections were defined as mild plus at least 1 of the following findings: increased C-reactive protein, leukocytosis, fever, or pus at the punctured site. Severe infections were defined by the need for a surgical incision or revision.

Selected variables for pairwise matching were sex, age, BMI, ASA physical status score, diabetes, anticoagulant therapy, antibiotic prophylaxis, catheter, single-shot blocks, nerve stimulation, department of surgery, sciatic nerve catheter (as second catheter), year of surgery, and hospital center.




FIGURE 3. Psoas block versus femoral block: ORs of complications. Odds ratios with 95% CI complications are displayed crude (A, B), adjusted for confounders (C, D), and before (A, C) and after matching (B, D). Risk parameters in C were adjusted as follows: vascular puncture was adjusted for ASA physical status score, diabetes, anticoagulant therapy, type of block, use of nerve stimulator, surgery, multiple skin punctures, year of surgery, and hospital. Multiple skin puncture variable was adjusted for age, BMI, ASA score, diabetes, type of block, use of nerve stimulator, surgery, year of surgery, and hospital. Primary failure was adjusted for ASA score, diabetes, anticoagulant therapy, use of nerve stimulator, year of surgery, and hospital. Infection any grade was adjusted for sex, BMI, ASA score, diabetes, antibiotic prophylaxis, type of block, surgery, catheter duration, year of surgery, and hospital. Patient and provider satisfaction was adjusted for sex, age, BMI, ASA score, diabetes, antibiotic prophylaxis, type of block, use of nerve stimulator, surgery, sciatic nerve block, vascular puncture, multiple skin punctures, primary failure of block, paresthesia during block insertion, block-related infection, catheter duration, year of surgery, and hospital. Risk parameters in D were adjusted for BMI, antibiotic prophylaxis, type of block, use of nerve stimulator, use of ultrasound, year of surgery, and hospital. Patient and provider satisfaction was additionally adjusted for vascular puncture, multiple skin punctures, primary failure of block, paresthesia during block insertion, block-related infection, and catheter duration.

sensitivity analysis. One strong aspect of our study is that block-related complications were clearly defined beforehand.

Our findings are generally consistent with previously reported risks of complications during insertion of femoral or sciatic nerve blocks: vascular puncture (present data: 1.1% vs Popping et al¹⁹: 4.4%), multiple skin punctures (present data: 7.7% vs Popping et al¹⁹: 9.1%), paresthesia (present data: 0.8% vs Popping et al¹⁹: 0.9%), and local anesthetic systemic toxicity (present data: 0% vs Popping et al¹⁹: 0%).¹⁹ However, our observed incidence of femoral catheter-related infections (0.9%) was lower than the 1.5% to 2.4% reported previously.^{10,19,20} Various definitions of infection and inflammation, duration of catheter use, preventive hygiene measures, and probably many unknown factors most likely led to these differences.

Auroy et al⁸ noted “80/10,000 serious complications” when psoas blocks were used. However, there were only 394 psoas blocks in their sample of voluntary reports, making their extrapolation highly unreliable. Another study reported no complications in 213 psoas blocks, which also provided an unreliable estimate.²¹ In contrast, our estimate of 0.3% in 6697 continuous psoas blocks has a much larger denominator and use compulsory documentation.

Our results are consistent with previous reports showing that risk is greater during insertion of psoas than femoral blocks. For

example, there are reports of fatal local anesthetic systemic toxicity and unnoticed accidental dural puncture.^{8,22} Anesthesiologists should be aware of these major complications and have been advised to manage the psoas block with the same vigilance as a neuraxial procedure.⁸ Moreover, anesthesiologists should be cognizant of arterial and venous anatomy within the lumbar nerve plexus and recognize that a psoas block can result in neuraxial anesthesia.⁸ Therefore, this block should be performed by well-trained physicians to avoid life-threatening complications during and after the procedure.²³ In our analysis of 7593 psoas blocks, no death was reported. Therefore, we assume that the psoas block is safe as previously reported.

Femoral blocks are easier to perform than psoas blocks. Consequently, premature termination of block attempts was less common with femoral blocks. However, only 0.1% of psoas block attempts in our series terminated early—a fraction that is so low that it does not provide a basis for selecting 1 type of block over the other. Once positioned, though, primary failure was more common with femoral (2.3%) than psoas (0.8%) blocks.

Satisfaction with analgesia for both providers and patients was higher in patients with psoas than femoral blocks. The reason could be a better quality of analgesia because of the included

obturator nerve.^{24,25} Moreover, block-related infections were far less common with psoas than femoral blocks (0.3% vs 0.9%; $P < 0.001$). Therefore, the psoas block seems to be a good block for lower-limb surgery.

Unfortunately, we cannot determine whether the observed, mostly minor, complications were linked with prolonged hospital stay or worse long-term outcomes because the duration of hospitalization, long-term outcomes, and long-term mortality were not recorded. The observed higher satisfaction in psoas block cannot be adequately assessed based on the results of this study. Residual confounding may introduce error, which will not be eliminated by either multivariable or propensity-matched analysis as in any nonrandomized analysis.

Our results were adjusted for the year of surgery, regardless of presumed improvements in knowledge, skills, techniques, and disinfectant methods during the 8-year observation period. The care site was considered a potential confounder in our statistical model, even though there was heterogeneity in the incidence of complications among the hospitals in our analysis. Statistically significant associations that are not clinically important could be identified because of our large patient population; however, the magnitude of the associations we observed was clearly clinically meaningful. Although our analysis was retrospective, complications data in our registry were specifically collected concurrent with patient care using a priori quality to ensure validity.

In conclusion, psoas blocks are associated with more insertion-related complications, but fewer infectious complications.

ACKNOWLEDGMENTS

The German Network for Regional Anesthesia investigators are all from participating hospital centers that collected data for the present study: Berlin—Charité CCM/CVK, Univ Prof Dr Claudia Spies; Berlin—DRK Kliniken Westend, Prof Dr Med Arnd Timmermann; Bochum—Knappschafts Krankenhaus, Prof Dr Michael Adamzik; Bad Saarow—HELIOS Klinikum, Dr Med Stefan Wirtz; Dresden—Uniklinikum, Prof Dr Med Thea Koch; Erfurt—HELIOS Klinikum, Prof Dr Med Andreas Meier-Hellmann and Dr Med Gerald Burgard; Frankfurt—Orthopädische Uniklinik, Prof Dr Med Paul Kessler; Frankfurt—BGU, Dr Med Rolf Teßmann; Hamburg—UKE, Univ Prof Dr Med Alwin E. Goetz; Hannover—Diakoniekrankenhaus Friederikenstift gGmbH, PD. Dr Med André Gottschalk; Jena—Uniklinikum, Prof Dr Med Konrad Reinhart; Lengerich—HELIOS Klinikum, Dr Med Albrecht Pfeiff; Ludwigsburg—Klinikum, Prof Dr Med Götz Geldner; Marburg—Uniklinikum, Prof Dr Med Hinnerk F. W. Wulf; Siegen—St Marien-Krankenhaus, Prof Dr Med Werner Hering; Städtisches Klinikum Solingen gGmbH, Prof Dr Med Thomas Standl; Ulm—Rehabilitationskrankenhaus, Dr Med Jörg Winkelmann.

REFERENCES

- Chayen D, Nathan H, Chayen M. The psoas compartment block. *Anesthesiology*. 1976;45:95–99.
- de Leeuw MA, Zuurmond WW, Perez RS. The psoas compartment block for hip surgery: the past, present, and future. *Anesthesiol Res Pract*. 2011;2011:159541.
- Aveline C, Bonnet F. Delayed retroperitoneal haematoma after failed lumbar plexus block. *Br J Anaesth*. 2004;93:589–591.
- Webster CS, Larsson L, Frampton CM, et al. Clinical assessment of a new anaesthetic drug administration system: a prospective, controlled, longitudinal incident monitoring study. *Anaesthesia*. 2010;65:490–499.
- Horlocker TT, Wedel DJ, Rowlingson JC, et al. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine evidence-based guidelines (third edition). *Reg Anesth Pain Med*. 2010;35:64–101.
- Capdevila X, Coimbra C, Choquet O. Approaches to the lumbar plexus: success, risks, and outcome. *Reg Anesth Pain Med*. 2005;30:150–162.
- Gentili M, Aveline C, Bonnet F. Total spinal anesthesia after posterior lumbar plexus block [in French]. *Ann Fr Anesth Reanim*. 1998;17:740–742.
- Auroy Y, Benhamou D, Bargues L, et al. Major complications of regional anesthesia in France: the SOS regional anesthesia hotline service. *Anesthesiology*. 2002;97:1274–1280.
- Barrington MJ, Kluger R. Ultrasound guidance reduces the risk of local anesthetic systemic toxicity following peripheral nerve blockade. *Reg Anesth Pain Med*. 2013;38:289–297.
- Capdevila X, Pirat P, Bringuier S, et al. Continuous peripheral nerve blocks in hospital wards after orthopedic surgery: a multicenter prospective analysis of the quality of postoperative analgesia and complications in 1,416 patients. *Anesthesiology*. 2005;103:1035–1045.
- Parienti JJ, Mongardon N, Megarbane B, et al. Intravascular complications of central venous catheterization by insertion site. *N Engl J Med*. 2015;373:1220–1229.
- Volk T, Engelhardt L, Spies C, et al. A German network for regional anaesthesia of the scientific working group regional anaesthesia within DGA and BDA [in German]. *Anesthesiol Intensivmed Notfallmed Schmerzther*. 2009;44:778–780.
- AdK-uPd A. Perioperative Antibiotikaprophylaxe. Registernummer: 029-022. Available at: www.awmf.org/leitlinien/detail/ll/029-022.html. Accessed March 21, 2016.
- Reutter F, Reuter DA, Hilgarth H, Heilek AM, Goepfert MS, Punke MA. Perioperative antibiotic prophylaxis [in German]. *Anaesthesist*. 2014;63:73–86.
- Morin AM, Kerwat KM, J. Büttner J, et al. Hygiene recommendations for the initiation and continued care of regional anaesthetic procedures—the 15 “musts” of the scientific working group regional anaesthesia. *Anesthesiol Intensivmed*. 2006;47:372–379.
- Büttner J, Buerkle H, Gogarten W, Wulf H. Thromboembolieprophylaxe bei peripheren Blockadetechniken zur Regionalanästhesie. *Anesth Intensivmed*. 2005;46:319–322.
- Bomberg H, Albert N, Schmitt K, et al. Obesity in regional anesthesia—a risk factor for peripheral catheter-related infections. *Acta Anaesthesiol Scand*. 2015;59:1038–1048.
- Bomberg H, Kubulus C, List F, et al. Diabetes: a risk factor for catheter-associated infections. *Reg Anesth Pain Med*. 2015;40:16–21.
- Popping DM, Zahn PK, van Aken HK, Dasch B, Boche R, Pogatzki-Zahn EM. Effectiveness and safety of postoperative pain management: a survey of 18 925 consecutive patients between 1998 and 2006 (2nd revision): a database analysis of prospectively raised data. *Br J Anaesth*. 2008;101:832–840.
- Cuvillon P, Ripart J, Lalourcey L, et al. The continuous femoral nerve block catheter for postoperative analgesia: bacterial colonization, infectious rate and adverse effects. *Anesth Analg*. 2001;93:1045–1049.
- Nye ZB, Horn JL, Crittenden W, Abrahams MS, Aziz MF. Ambulatory continuous posterior lumbar plexus blocks following hip arthroscopy: a review of 213 cases. *J Clin Anesth*. 2013;25:268–274.
- Vadi MG, Patel N, Stiegler MP. Local anesthetic systemic toxicity after combined psoas compartment-sciatic nerve block: analysis of decision factors and diagnostic delay. *Anesthesiology*. 2014;120:987–996.
- Macaire PGE, Choquet O. Le bloc du plexus lombaire est-il dangereux? In: *Évaluation et Traitement de la Douleur SFAR 2002*. Paris, France: Elsevier et SFAR Ed; 2002:37–50.
- Bendtsen TF, Moriggl B, Chan V, Borglum J. The optimal analgesic block for total knee arthroplasty. *Reg Anesth Pain Med*. 2016;41:711–719.
- Runge C, Borglum J, Jensen JM, et al. The analgesic effect of obturator nerve block added to a femoral triangle block after total knee arthroplasty: a randomized controlled trial. *Reg Anesth Pain Med*. 2016;41:445–451.